HIERARCHICAL DECOMPOSITION OF DOMAINS WITH FRACTURES

SUSANNA GEBAUER, RALF KORNHUBER, AND HARRY YSERENTANT

ABSTRACT. We consider the efficient and robust numerical solution of elliptic problems with jumping coefficients occurring on a network of thin fractures. We present an iterative solution concept based on a hierarchical separation of the fractures and the surrounding rock matrix. Upper estimates for the convergence rates are independent of the width of the fractures and of the jumps of the coefficients. Inexact solution of the local subproblems is also considered. The theoretical results are illustrated by numerical experiments.

1. INTRODUCTION

Saturated groundwater flow in fractured porous media can be described by linear elliptic problems. Fractures on one or more micro scales are usually represented by effective parameters of corresponding single or multi-porosity models based on homogenization techniques. Fractures on macro scales directly enter the geometry of the mathematical model. For lack of data, such fracture networks are typically generated automatically based on stochastic reasoning [20]. The permeability k_F within the fractures is usually some orders of magnitude larger than the permeability of the surrounding rock matrix, while their width ϵ_F might be some orders of magnitude smaller than the overall computational domain. Similar problems occur in other applications: The heat transfer in the human body is dominated by the blood vessels which, depending on their size, are represented by effective parameters [7, 8] or have to be incorporated directly. Another example concerns diffusion induced drug permeation through stratum corneum [11, 16]. In this case, the lipid "mortar" between the corneocytes plays the role of the fractures.

In order to avoid numerical troubles resulting from small width ϵ_F and large permeability k_F, fractures (or vessels) are often discretized by lower dimensional elements [2, 12, 13] (or [18]). However, there are also some disadvantages of this approach. For example, outward normal flow and mass conservation across the interface are not incorporated. This motivated recent work on equidimensional discretizations [9, 10, 14, 15].

On this background, we consider the efficient and robust numerical solution of elliptic problems with jumping coefficients occurring on a network of long, thin fractures. Robustness means that the complexity should not depend on the crucial parameters ϵ_F and k_F. To this end, the fracture network is discretized by anisotropic isoparametric bilinear finite elements while usual triangular elements are used elsewhere. The main part of the paper concentrates on the iterative solution of the resulting discrete problems. We introduce so-called *hierarchical domain*

This work has been funded in part by the German Science foundation under contract Ko 1806/2-1 and Ko 1806/2-3 .

decomposition methods (HDD) based on a hierarchical splitting of the discrete solution space into three subspaces associated with the interior of the fractures, the interface, and the surrounding rock matrix. The matrix space consists of functions which are essentially constant across the fractures. We show that the stability of this decomposition and therefore the convergence speed of the resulting subspace correction method is independent of ϵ_F, k_F and of the mesh size. The proof is based on the assumption that the fractures are "almost" lower dimensional in the sense that their resolution by isotropic elements is not desirable, e.g., for efficiency reasons. This setting allows to trace back the stability of the hierarchical decomposition to the optimal stability properties of one-dimensional interpolation. As a consequence, our approach is applicable in any space dimension. Exceptional situations like crossings of fractures or fractures ending in the computational domain can be resolved by local estimates on the surrounding elements. The exact solution of the three local subproblems can be replaced by suitable multigrid methods. We present an illustrative example adapting well-known results on hierarchical bases. In this case, the number of refinement steps enters only polynomially and independence of ϵ_F and, in particular, of k_F is preserved. However, this result is restricted to two space dimensions. In our numerical experiments, we observe similar convergence speed as for classical multigrid methods applied to the Laplace equation.

In a sense, our approach is complementary to the algorithm presented by Heisig et al. [11] which aims at compensating small ε_F by successive anisotropic refinement and accounts for large k_F by ILU-smoothing. Algebraic multigrid methods (see, e.g., [4, 19] or [22] for an overview) usually work reasonably well but mostly suffer from a certain lack of theory. Apel and Schöberl [1] consider a completely different anisotropic problem with a tensor product structure, where line (or plane) smoothers or semi-coarsening can be applied.

The paper is organized as follows. We first present a model problem with two fractures crossing in a two-dimensional domain, its discretization and an a priori estimate of the discretization error which is robust with respect to ϵ_F and k_F. The next two sections concentrate on the stable separation of the subspace associated with the interior of the fractures and the interface. In Section 5 we present the resulting HDD algorithm together with an inexact variant and derive upper bounds for the convergence rates. Numerical experiments confirm our theoretical findings.

2. A discrete elliptic problem on a domain with fractures

Let $\Omega \subset \mathbb{R}^2$ be a polygonal domain, e.g. the unit square, with the two fractures

$$\Omega_F^i = \{x \in \Omega \mid x = b_F^i + s d_F^i + t n_F^i,\ s \in \mathbb{R},\ t \in (0, \varepsilon_F)\}, \quad i = 1, 2,$$

each of which is characterized by its position vector b_F^i, direction d_F^i, normal n_F^i and width $\epsilon_F > 0$. We assume that the fractures cross inside of Ω, i.e. that $\overline{\Omega}_c \subset \Omega$, $\Omega_c = \Omega_F^1 \cap \Omega_F^2$. The network of fractures is denoted by $\Omega_F = \Omega_F^1 \cup \Omega_F^2$ while $\Omega_M = \Omega \setminus \overline{\Omega}_F$ and $\Gamma = \overline{\Omega}_M \cap \overline{\Omega}_F$ represent the rock matrix and the interface, respectively. This leads to the decomposition

$$\Omega = \Omega_F \cup \Gamma \cup \Omega_M. \tag{2.1}$$

We consider the elliptic variational problem

$$u \in H : \qquad a(u, v) = \ell(v) \qquad \forall v \in H \tag{2.2}$$

with the symmetric bilinear form $a(v, w) = (K\nabla u, \nabla v)_{L^2(\Omega)}$ and jumping permeability K,

$$K(x) = \begin{cases} k_F \geq 1, & x \in \Omega_F \\ 1, & x \in \Omega_M \end{cases} . \tag{2.3}$$

For simplicity, let $H = H_0^1(\Omega)$ and $\ell \in H'$ is some right hand side. The energy norm is denoted by $\|\cdot\| = a(\cdot,\cdot)^{1/2}$.

Let $\mathcal{P}_0 = \mathcal{T}_0 \cup \mathcal{Q}_0$ be a subdivision of $\Omega = \overline{\Omega}_M \cup \overline{\Omega}_F$ consisting of the partitions

$$\overline{\Omega}_M = \bigcup_{T \in \mathcal{T}_0} T, \qquad \overline{\Omega}_F = \bigcup_{Q \in \mathcal{Q}_0} Q$$

into triangles T and trapezoidals Q, respectively. We assume that the vertices of each trapezoidal $Q \in \mathcal{Q}_0$ lie on Γ (see the left picture in Figure 2.1). In particular,

$$Q_c := \overline{\Omega}_c \in \mathcal{Q}_0$$

is a parallelogram. We further assume that $\mathcal{P}_0$ is *conforming* in the sense that the intersection of two different elements is either a common edge, a common vertex or empty. Finally, $\mathcal{P}_0$ is supposed to be *shape regular* in the sense that all $P \in \mathcal{P}_0$ have positive area and all $Q \in \mathcal{Q}_0$ have four different vertices. Equivalently, there are positive constants s_0, $\gamma_0 \in \mathbb{R}$ such that

$$s_0 h_Q \leq h'_Q, \quad \forall Q \in \mathcal{Q}_0 \setminus \{Q_c\}, \qquad \gamma_0 \leq \gamma_P \leq \pi - \gamma_0 \quad \forall P \in \mathcal{P}_0 \tag{2.4}$$

holds with $h_Q \geq h'_Q$ denoting the lengths of the two parallel edges contained in Γ and γ_P is an arbitrary interior angle. Note that all edges of Q_c have the same length h_{Q_c}.

Though all results and algorithms to be presented can be directly extended to locally refined grids, we assume for simplicity that the triangulation $\mathcal{T}_1$ is obtained by uniform refinement of $\mathcal{T}_0$. More precisely, each triangle $T \in \mathcal{T}_0$ is subdivided into four similar subtriangles. Connecting the resulting new midpoints of opposite edges of each $Q \in \mathcal{Q}_0 \setminus \{Q_c\}$, we get the set $\mathcal{Q}_1$ of anisotropically refined trapezoidals . Reiteration of this procedure leads to a sequence of refined partitions $\mathcal{P}_j = \mathcal{T}_j \cup \mathcal{Q}_j$, $j = 0, 1, \ldots$. Observe that $\mathcal{P}_j$ is conforming and the shape regularity (2.4) is preserved uniformly in j. In particular, the angle condition in (2.4) implies

$$h_Q - h'_Q \leq \frac{\varepsilon_F}{(\sin\gamma_0)^2} = \mathcal{O}(\varepsilon_F) \qquad \forall Q \in \mathcal{Q}_j \tag{2.5}$$

so that the elements of $\mathcal{Q}_j$ tend to (highly anisotropic) parallelograms as ε_F tends to zero.

We assume that the *fractures are long, thin objects* in the sense that the resolution of the corresponding anisotropy of the elements $Q \in \mathcal{Q}_0$ by the above refinement procedure is not desirable, e.g., for efficiency reasons. More precisely, we impose the condition

$$\varepsilon_F \leq C_0 h_Q \qquad \forall Q \in \mathcal{Q}_j \tag{2.6}$$

that the width of the fractures is bounded by the size of the surrounding mesh.

Anisotropic refinement of $\mathcal{Q}_j$ in the other direction is performed by bisecting and connecting the midpoints of all edges not contained in Γ. Application of k steps of this procedure to $\mathcal{P}_j$ provides the partition $\mathcal{P}_{jk} = \mathcal{T}_j \cup \mathcal{Q}_{jk}$ (see the right picture of Figure 2.1 for $j = 1$, $k = 2$). The refined partitions $\mathcal{P}_{jk}$ are conforming and the



FIGURE 2.1. Initial partition $\mathcal{P}_0$ and refined partition $\mathcal{P}_{12}$

shape regularity (2.4) holds uniformly in $j, k \in \mathbb{N}$. Finally note that $\mathcal{P}_{jk}$ does not depend on the order of the above two types of refinement steps.

For given j, k, let $\mathcal{S}_{jk} \in H_0^1(\Omega)$ be the subspace of functions v such that $v|_T$ is linear and $v|_Q$ is isoparametric bilinear for all $T \in \mathcal{T}_j$ and $Q \in \mathcal{Q}_{jk}$, respectively. Then the corresponding finite element discretization of the continuous problem (2.2) reads as follows

$$u_{jk} \in \mathcal{S}_{jk} : \qquad a(u_{jk}, v) = \ell(v) \qquad \forall v \in \mathcal{S}_{jk}. \tag{2.7}$$

In preparation for an error estimate, we introduce the weighted Sobolev norms

$$\|v\|_{m,K} = \sum_{|\alpha| \le m} \int_\Omega K(x)(D^\alpha v(x))^2 \, dx, \tag{2.8}$$

for $m = 0, 1, \ldots$, using standard multi-index notation (cf., e.g. [6, Chapter 1]). The obvious norm equivalence

$$\|v\|_{H^m(\Omega)} \le \|v\|_{m,K} \le k_F \|v\|_{H^m(\Omega)}$$

directly extends to the corresponding intermediate norms $\|v\|_{H^s(\Omega)}$ $\|v\|_{s,K}$, $s \in \mathbb{R}$, as obtained by interpolation (cf. Bergh and Löfström [3] or Brenner and Scott [6, Chapter 12]).

Proposition 2.1. *Assume that $u \in H^{1+s}(\Omega)$ with $0 < s \le 1$. Then the finite element solution u_{jk} satisfies the error estimate*

$$\|u - u_{jk}\| \le C h_{jk}^s \|u\|_{1+s,K}, \qquad h_{jk} = \max_{P \in P_{jk}} \operatorname{diam} P, \tag{2.9}$$

with a constant $C = C(s_0, \gamma_0)$ depending only on the shape regularity (2.4) of $\mathcal{P}_0$.

Proof. Utilizing the Lax-Milgram lemma together with standard estimates of the interpolation error on isotropic triangles or trapezoidals and the results of Ženíšek and Vanmaele [21] for the anisotropic case, we obtain the estimate

$$\|u - u_{jk}\| \le c h_{jk} \|u\|_{2,K}$$

provided that $u \in H^2(\Omega)$. Here, the constant c depends only on the shape regularity of $P \in \mathcal{P}_{jk}$ and therefore of $\mathcal{P}_0$. Now the desired estimate (2.9) follows from standard results on interpolated Sobolev spaces [3],[6, Chapter 12]. ☐

3. Separation of the fractures

We consider the direct splitting

$$\mathcal{S}_{jk} = \mathcal{S}_j^{\overline{M}} \oplus \mathcal{S}_{jk}^F \tag{3.1}$$

of the finite element space $\mathcal{S}_{jk}$ into the fracture space

$$\mathcal{S}_{jk}^F = \left\{ v \in \mathcal{S}_{jk} \mid v|_{\overline{\Omega}_M} = 0 \right\} \tag{3.2}$$

and its complement $\mathcal{S}_j^{\overline{M}}$ consisting of all $v \in \mathcal{S}_{jk}$ such that $v|_Q$ is isoparametric bilinear for all $Q \in \mathcal{Q}_j$. The stability of the splitting (3.1) is equivalent to the stability of the interpolation operator $I_{jk}^{\overline{M}} : \mathcal{S}_{jk} \to \mathcal{S}_j^{\overline{M}}$ defined by

$$I_{jk}^{\overline{M}} v(p) = v(p), \qquad p \in \mathcal{N}_j^{\overline{M}}.$$

Here, $\mathcal{N}_j^{\overline{M}}$ denotes the set of all the vertices of $P \in \mathcal{P}_{jk}$ which lie in $\overline{\Omega}_M$. Stability of $I_{jk}^{\overline{M}}$ will be shown by local estimates. To this end, we first consider transformations to suitable reference elements.

The nodes of $Q = (p_1, \ldots, p_4) \in \mathcal{Q}_j$ are ordered anti-clockwise and, for $Q \neq Q_c$, in such a way that the lengths h_Q, h_Q' of the edges $[p_1, p_2]$, $[p_3, p_4]$ lying in Γ satisfy $h_Q' \leq h_Q$. With each $Q \neq Q_c$ we associate the reference element $\hat{Q} = [0,1] \times [0, \varepsilon]$ where $\varepsilon = \varepsilon_F / h_Q$. Observe, that we always have $\varepsilon \leq C_0$ by (2.6), but $\varepsilon \to 0$ occurs for $\varepsilon_F \to 0$. For the isotropic crossing Q_c we select the canonical reference element $\hat{Q}_c = [0,1]^2$. Let $\mathcal{F}_Q : \hat{Q} \to Q$ denote the bijective bilinear mapping such that the ordering of the vertices is preserved and $\mathcal{F}_Q(0,0) = p_1$.

Throughout this paper, we write $a \preceq b$ for $a \leq Cb$ and $a \asymp b$ for $cb \leq a \leq Cb$ with some c, C depending only on the constants s_0, γ_0 and C_0 from (2.4) and (2.6), respectively.

Lemma 3.1. *Let $Q \in \mathcal{Q}_j$. Each $v \in H^1(Q)$ and its transformation $\hat{v}$,*

$$\hat{v} = v(\mathcal{F}_Q(\cdot)) : \hat{Q} \to \mathbb{R},$$

satisfy the norm equivalence

$$\|\nabla v\|_{L^2(Q)} \asymp \|\nabla \hat{v}\|_{L^2(\hat{Q})}. \tag{3.3}$$

Proof. As the assertion is well-known for $Q = Q_c$, we only have to consider the remaining case $Q \neq Q_c$. The chain rule yields

$$\nabla \hat{v} = \nabla v B_Q$$

denoting $B_Q = \mathcal{F}_Q'$. Elementary calculations utilizing (2.4) and (2.5) provide

$$|B_Q| \preceq (1 + \tfrac{h_Q - h_Q'}{\varepsilon_F}) \max\{h_Q, \frac{\varepsilon_F}{\varepsilon}\} = h_Q$$

where $|\cdot|$ is the spectral norm. Similarly, we get

$$|B_Q^{-1}| \preceq (1 + \tfrac{h_Q - h_Q'}{\varepsilon_F}) \max\{h_Q^{-1}, \frac{\varepsilon}{\varepsilon_F}\} \preceq h_Q^{-1}$$

and in addition

$$|\det B_Q| \leq h_Q \frac{\varepsilon_F}{\varepsilon} = h_Q^2, \quad |\det B_Q^{-1}| \preceq h_Q^{-1} \frac{\varepsilon}{\varepsilon_F} = h_Q^{-2}$$

so that the assertion follows from the substitution rule for integrals. □

The following technical lemma allows to extend basic arguments from one-dimensional intervals to long, thin elements in higher dimensions. In this sense, it is crucial for the rest of this paper.

Lemma 3.2. *Let* $w : \hat{Q} = [0,1] \times [0,\varepsilon] \to \mathbb{R}$ *be such that* $w(\xi,\cdot)$ *is absolutely continuous for each fixed* $\xi \in [0,1]$*,* $w(\cdot,\eta)$ *is linear for each fixed* $\eta \in [0,\varepsilon]$*, and* $w(\xi,0) = w(\xi,\varepsilon) = 0$ *holds for all* $\xi \in [0,1]$*. Then*

$$\|\nabla w\|^2_{L^2(\hat{Q})} \le (1 + 12\varepsilon^2)\|\tfrac{\partial}{\partial\eta} w\|^2_{L^2(\hat{Q})}. \tag{3.4}$$

Proof. The assertion will be proved in three steps. First, let $f : [0,\varepsilon] \to \mathbb{R}$ be absolutely continuous and $f(0) = 0$. Then the fundamental theorem of calculus together with Cauchy's inequality yields

$$\int_0^\varepsilon (f(\eta))^2 \, d\eta = \int_0^\varepsilon \left(\int_0^s 1 \cdot f'(s) \, ds \right)^2 d\eta \le \varepsilon^2 \int_0^\varepsilon (f'(s))^2 \, ds. \tag{3.5}$$

Next, let $g : [0,1] \to \mathbb{R}$ be linear. Then elementary calculation leads to

$$(g(1) - g(0))^2 \le 12 \int_0^1 (g(\xi))^2 \, d\xi \tag{3.6}$$

where, in particular, the binomial estimate

$$(a-b)^2 \le 4(a^2 + ab + b^2) \qquad \forall a,b \in \mathbb{R}$$

has been used. Inserting $f(\eta) = w(1,\eta) - w(0,\eta)$ and $g(\xi) = \frac{\partial}{\partial\eta} w(\xi,\eta)$ for fixed η, in (3.5) and (3.6), respectively, the assertion follows from

$$\begin{aligned}\int_0^\varepsilon \int_0^1 \left(\tfrac{\partial}{\partial\xi} w(\xi,\eta)\right)^2 d\xi \, d\eta &= \int_0^\varepsilon (w(1,\eta) - w(0,\eta))^2 \, d\eta \\ &\le \varepsilon^2 \int_0^\varepsilon \left(\tfrac{\partial}{\partial\eta}(w(1,\eta) - w(0,\eta))\right)^2 d\eta \le 12\varepsilon^2 \int_0^\varepsilon \int_0^1 \left(\tfrac{\partial}{\partial\eta} w(\xi,\eta)\right)^2 d\xi \, d\eta.\end{aligned}$$

□

Now we can state a local estimate for anisotropic elements $Q \neq Q_c$.

Lemma 3.3. *The estimate*

$$\|\nabla I^{\overline{M}}_{jk} v\|_{L^2(Q)} \preceq \|\nabla v\|_{L^2(Q)} \tag{3.7}$$

holds for all $v \in \mathcal{S}_{jk}$ *and* $Q \in \mathcal{Q}_j \setminus \{Q_c\}$*.*

Proof. Transformation of $I^{\overline{M}}_{jk} v$ to the reference element $\hat{Q} = [0,1] \times [0,\varepsilon]$ provides $\hat{I}\hat{v}$, where $\hat{v}$ is the transformation of some $v \in \mathcal{S}_{jk}$ and $\hat{I}$ denotes the bilinear interpolation at the vertices of $\hat{Q}$. It is easily checked that $w = \hat{v} - \hat{I}\hat{v}$ satisfies the assumptions of Lemma 3.2. Hence,

$$\|\nabla(\hat{v} - \hat{I}\hat{v})\|^2_{L^2(\hat{Q})} \le (1 + 12\varepsilon^2)\|\tfrac{\partial}{\partial\eta}(\hat{v} - \hat{I}\hat{v})\|^2_{L^2(\hat{Q})}.$$

Using the orthogonality

$$\int_0^\varepsilon \tfrac{\partial}{\partial\eta}(\hat{v} - \hat{I}\hat{v}) \tfrac{\partial}{\partial\eta} \hat{I}\hat{v} \, d\eta = 0$$

we get

$$\|\tfrac{\partial}{\partial\eta}(\hat{v} - \hat{I}\hat{v})\|^2_{L^2(\hat{Q})} \le \|\tfrac{\partial}{\partial\eta}\hat{v}\|^2_{L^2(\hat{Q})}$$

so that the assertion follows from Lemma 3.1. □

The proof of Lemma 3.3 extends the stability of nodal interpolation from intervals to long, thin elements in two space dimensions. We emphasize that stability of nodal interpolation on anisotropic elements in three space dimensions can be derived in a similar way.

The above technique cannot be applied to the isotropic crossing Q_c. The basic idea to treat such exceptional elements is first to estimate the norm of $I_{jk}^{\overline{M}} v$ on Q_c by the norm of $I_{jk}^{\overline{M}} v$ on a neighboring element $Q \in \mathcal{Q}_j \setminus Q_c$ and then to apply the preceding Lemma 3.3. Observe that $v|_{Q_c}$ does not enter the resulting upper bound.

Lemma 3.4. *There is a neighboring element $Q \in \mathcal{Q}_j$ of Q_c such that the estimate*

$$\|\nabla I_{jk}^{\overline{M}} v\|_{L^2(Q_c)} \preceq \|\nabla v\|_{L^2(Q)} \tag{3.8}$$

holds for all $v \in \mathcal{S}_{jk}$.

Proof. Let $\hat{w}$ be a bilinear function on $\hat{Q} = [0,1] \times [0,\varepsilon]$ interpolating the values $w_1, \ldots, w_4$ in the vertices $(0,0)$, $(1,0)$, $(1,\varepsilon)$, $(0,\varepsilon)$. Elementary calculation yields

$$\begin{aligned} \|\nabla \hat{w}\|^2_{L^2(\hat{Q})} = \tfrac{1}{3}\varepsilon \left((w_2 - w_1)^2 + (w_3 - w_4)^2 + (w_2 - w_1)(w_3 - w_4)\right) \\ + \tfrac{1}{3}\varepsilon^{-1} \left((w_4 - w_1)^2 + (w_3 - w_2)^2 + (w_4 - w_1)(w_3 - w_2)\right). \end{aligned} \tag{3.9}$$

Hence, the binomial estimate

$$\tfrac{4}{3}(a^2 + b^2 + ab) \geq a^2 \qquad \forall a, b \in \mathbb{R} \tag{3.10}$$

provides

$$4\varepsilon \|\nabla \hat{w}\|^2_{L^2(Q_c)} \geq \max\{(w_4 - w_1)^2, (w_3 - w_2)^2\}. \tag{3.11}$$

Replacing $\hat{Q}$ by $\hat{Q}_c = [0,1]^2$, formula (3.9) obviously holds with $\varepsilon = 1$. Using $ab \leq \frac{1}{2}(a^2 + b^2)$ we easily get

$$\|\nabla \hat{w}\|^2_{L^2(Q_c)} \leq 2 \max_{l=1,\ldots,4} (w_{l+1} - w_l)^2 \tag{3.12}$$

denoting $w_5 = w_1$.

Now let $v \in \mathcal{S}_{jk}$ and $w = I_{jk}^{\overline{M}} v$. Using (3.12) together with Lemma 3.1 we obtain without loss of generality

$$\|\nabla w\|^2_{L^2(Q_c)} \preceq \max_{l=1,\ldots,4} (w_{l+1} - w_l)^2 = (w_4 - w_1)^2.$$

In this case, we select $Q = (p_1, p_2, p_3, p_4) \in \mathcal{Q}_j$ of Q_c such that $p_1, p_4 \in Q \cap Q_c$. Then (3.11) and Lemma 3.1 provide the estimate

$$\|\nabla w\|_{L^2(Q_c)} \preceq \|\nabla w\|_{L^2(Q)}$$

and the assertion follows from the preceding Lemma 3.3. □

Estimates of the norm of $I_{jk}^{\overline{M}} v$ on other exceptional elements, occurring, e.g., at the end of a fracture, by its norm on neighboring elements can be derived in a similar way. We also refer to the proof of Proposition 4.1 later on.

We are ready for the main result of this section.

Proposition 3.1. *For each $v \in \mathcal{S}_{jk}$ the decomposition $v = v^F + v^{\overline{M}}$ into $v^F \in \mathcal{S}_{jk}^F$ and $v^{\overline{M}} \in \mathcal{S}_j^{\overline{M}}$ satisfies*

$$\|v^F\|^2 + \|v^{\overline{M}}\|^2 \preceq \|v\|^2. \tag{3.13}$$

Proof. As a consequence of Lemma 3.3 and Lemma 3.4 we obtain

$$\|I_{jk}^{\overline{M}}v\|^2 = \sum_{T\in\mathcal{T}_j} \|\nabla v\|^2_{L^2(T)} + \sum_{Q\in\mathcal{Q}_j} k_F \|\nabla I_{jk}^{\overline{M}}v\|^2_{L^2(Q)} \preceq \|v\|^2$$

so that the assertion follows with $v^{\overline{M}} = I_{jk}^{\overline{M}}v$ and $v^F = v - I_{jk}^{\overline{M}}v$. □

The gradient of a function $v \in \mathcal{S}_j^{\overline{M}}$ with fixed nodal values typically becomes arbitrary large as $\varepsilon_F \to 0$. This causes problems with the robustness of iterative solvers for local subproblems on $\mathcal{S}_j^{\overline{M}}$ with respect to small ε_F. As a remedy, we now consider a further splitting of $\mathcal{S}_j^{\overline{M}}$ into a space of functions living on the interface Γ and another space of functions which are essentially constant across the fractures.

4. Separation of the interface

We decompose the interface $\Gamma = \Gamma_0 \cup \Gamma_1$ into its 'lower' and 'upper' part consisting of $\Gamma_l = \{x \in \Omega \mid x = b_F^i + sd_F^i + ln_F^i, s \in \mathbb{R}, i = 1, 2\}$ with $l = 0$ and $l = 1$, respectively. Let $\mathcal{N}_j^{\Gamma_0} = \mathcal{N}_j \cap \Gamma_0$ and $\mathcal{E}_j^F = \mathcal{E}_j \cap \overline{\Omega}^F$ with $\mathcal{N}_j$ and $\mathcal{E}_j$ denoting the sets of interior vertices and edges of $P \in \mathcal{P}_j$, respectively. We consider the direct splitting

$$\mathcal{S}_j^{\overline{M}} = \mathcal{S}_j^M \oplus \mathcal{S}_j^{\Gamma_0} \tag{4.1}$$

into the interface space

$$\mathcal{S}_j^{\Gamma_0} = \left\{ v \in \mathcal{S}_j^{\overline{M}} \mid v(p) = 0 \quad \forall p \notin \mathcal{N}_j^{\Gamma_0} \right\}, \tag{4.2}$$

and its complement $\mathcal{S}_j^M$ consisting of all $v \in \mathcal{S}_j^{\overline{M}}$ such that $v|_E$ is constant for all edges $E \in \mathcal{E}_j^F$. The splitting (4.1) is induced by the interpolation operator $I_j^M : \mathcal{S}_j^{\overline{M}} \to \mathcal{S}_j^M$ defined by

$$I_j^M v(p) = v(p) \qquad p \in \mathcal{N}_j \setminus \mathcal{N}_j^{\Gamma_0}.$$

If $p \in \mathcal{N}_j^{\Gamma_0}$, then $I_j^M v(p) = v(p^*)$ where p^* is the vertex of the edge $E = (p, p^*) \in \mathcal{E}_j^F$ or the vertex of Q_c which is not contained in $\mathcal{N}_j^{\Gamma_0}$. In particular, $I_j^M v$ is constant on Q_c.

We proceed with local stability estimates on $T \in \mathcal{T}_j$ and $Q \in \mathcal{Q}_j$, respectively.

Lemma 4.1. *The estimate*

$$\|\nabla I_j^M v\|^2_{L^2(T)} \preceq \|\nabla v\|^2_{L^2(T)} + \sum_{i=1}^{3} \left(I_j^M v(p_i) - v(p_i)\right)^2 \tag{4.3}$$

holds for all $v \in \mathcal{S}_j^{\overline{M}}$ and all $T = (p_1, \ldots, p_3) \in \mathcal{T}_j$.

Proof. We set $w = I_j^M v$ and $\hat{w}$, $\hat{v}$ denote the usual transformations of w, v to the reference element $T_0 = ((0,0), (1,0), (0,1))$. Elementary calculations provide

$$\|\nabla \hat{w}\|^2_{L^2(T_0)} = \tfrac{1}{2}\left((w_2 - w_1)^2 + (w_3 - w_1)^2\right) \le 2\|\nabla \hat{v}\|^2_{L^2(T_0)} + 4\textstyle\sum_{i=1}^{3}(w_i - v_i)^2$$

and the assertion follows from the well-known estimates $\|\nabla w\|_{L^2(T)} \preceq \|\nabla \hat{w}\|_{L^2(T_0)}$ and $\|\nabla \hat{v}\|_{L^2(T_0)} \preceq \|\nabla v\|_{L^2(T)}$. □

Lemma 4.2. *The estimates*

$$\|\nabla I_j^M v\|_{L^2(Q)} \preceq \|\nabla v\|_{L^2(Q)}, \qquad \max_{i=1,\dots,4} \big(I_j^M v(p_i) - v(p_i)\big)^2 \preceq \|\nabla v\|^2_{L^2(Q)} \tag{4.4}$$

hold for all $v \in \mathcal{S}_j^{\overline{M}}$ *and all* $Q = (p_1, \dots, p_4) \in \mathcal{Q}_j$.

Proof. We set $w = I_j^M v$ and $\hat{w}$, $\hat{v}$ denote the transformed functions on the reference element $\hat{Q} = [0,1] \times [0, \varepsilon]$. In the light of Lemma 3.1 it is sufficient to show

$$\|\nabla \hat{w}\|_{L^2(\hat{Q})} \preceq \|\nabla \hat{v}\|_{L^2(\hat{Q})}, \qquad \max_{i=1,\dots,4} (w_i - v_i)^2 \preceq \|\nabla \hat{v}\|^2_{L^2(\hat{Q})}, \tag{4.5}$$

respectively. Here, we have set $w_i = w(p_i)$, $v_i = v(p_i)$. As $\hat{v}$ is bilinear on $\hat{Q}$, we can use (3.11) to obtain

$$4\varepsilon \|\nabla \hat{v}\|^2_{L^2(\hat{Q})} \geq \max\{(v_4 - v_1)^2, (v_3 - v_2)^2\} = \max_{i=1,\dots,4} (w_i - v_i)^2$$

which is the right estimate in (4.4). Note that

$$\|\nabla \hat{w}\|^2_{L^2(\hat{Q})} = \varepsilon (w_2 - w_1)^2 \leq \varepsilon \max_{i,l=1,\dots,4} (v_i - v_l)^2 \leq 4\varepsilon \max_{l=1,\dots,4} (v_{l+1} - v_l)^2,$$

where we have set $v_5 = v_1$. Using the binomial estimate (3.10) and the representation (3.9) of $\|\nabla \hat{v}\|^2_{L^2(\hat{Q})}$ we further get

$$\varepsilon \max_{l=1,\dots,4} (v_{l+1} - v_l)^2 \leq 4(1 + \varepsilon^2) \|\nabla \hat{v}\|^2_{L^2(\hat{Q})}$$

which proves the assertion for $Q \neq Q_c$. In the case $Q = Q_c$ the left estimate in (4.4) is trivial and the right one follows by similar arguments as before. □

Now we are ready to prove stability of the splitting (4.1).

Proposition 4.1. *For each* $v \in \mathcal{S}_j^{\overline{M}}$ *the decomposition* $v = v^M + v^{\Gamma_0}$ *into* $v^M \in \mathcal{S}_j^M$ *and* $v^{\Gamma_0} \in \mathcal{S}_j^{\Gamma_0}$ *satisfies*

$$\|v^M\|^2 + \|v^{\Gamma_0}\|^2 \preceq \|v\|^2. \tag{4.6}$$

Proof. Let $v \in \mathcal{S}_j^{\overline{M}}$. We set $v^M = I_j^M v$ and $v_0^\Gamma = v - I_j^M v$. Utilizing Lemma 4.1 and Lemma 4.2 we get

$$\|\nabla v^M\|^2_{L^2(T)} \preceq \|\nabla v\|^2_{L^2(T)} + \sum_{Q \in \mathcal{Q}_j(T)} \|\nabla v\|^2_{L^2(Q)} \qquad \forall T \in \mathcal{T}_j$$

denoting $\mathcal{Q}_j(T) = \{Q \in \mathcal{Q}_j \mid Q \cap T \neq \emptyset\}$. As a consequence of the minimal angle condition (2.4) we have

$$\sum_{T \in \mathcal{T}_j} \sum_{Q \in \mathcal{Q}_j(T)} \|\nabla v\|^2_{L^2(Q)} \preceq \sum_{Q \in \mathcal{Q}_j} \|\nabla v\|^2_{L^2(Q)}.$$

Together with $k_F \geq 1$ and Lemma 4.2 these estimates provide

$$\|v^M\|^2 = \sum_{T \in \mathcal{T}_j} \|\nabla v^M\|^2_{L^2(T)} + \sum_{T \in \mathcal{Q}_j} k_F \|\nabla v^M\|^2_{L^2(Q)} \preceq \|v\|^2.$$

The assertion now follows from the triangle inequality. □

Note that Proposition 4.1 implies the stability of the overlapping splitting

$$\mathcal{S}_j^{\overline{M}} = \mathcal{S}_j^M + \mathcal{S}_j^\Gamma \tag{4.7}$$

where $\mathcal{S}_j^{\Gamma_0}$ is replaced by the larger space

$$\mathcal{S}_j^\Gamma = \left\{ v \in \mathcal{S}_j^{\overline{M}} \mid v(p) = 0 \quad \forall p \in \mathcal{N}_j \cap \Omega_M \right\}. \tag{4.8}$$

Utilizing Proposition 3.1, we also get the stability of the decomposition

$$\mathcal{S}_{jk} = \mathcal{S}_j^M + \mathcal{S}_{jk}^{\overline{F}}$$

denoting $\mathcal{S}_{jk}^{\overline{F}} = \mathcal{S}_{jk}^F + \mathcal{S}_j^\Gamma$.

5. Hierarchical domain decomposition methods

The successive subspace correction method (cf. Xu [23] or Yserentant [25]) resulting from a decomposition

$$\mathcal{S}_{jk} = \mathcal{W}_0 + \mathcal{W}_1 + \cdots + \mathcal{W}_J, \quad \mathcal{W}_l \subset \mathcal{S}_{jk},$$

and symmetric, positive definite bilinear forms $b_l(\cdot,\cdot)$ on $\mathcal{W}_l$ reads as follows: Starting with some given iterate $w_{-1} = u_{jk}^\nu \in \mathcal{S}_{jk}$, a sequence of intermediate iterates w_l, $l = 0, \dots, J$, is computed according to

$$\begin{aligned} v_l \in \mathcal{W}_l: \qquad b_l(v_l, v) &= \ell(v) - a(w_{l-1}, v) \qquad \forall v \in \mathcal{W}_l \\ w_{l+1} &= w_l + v_l \end{aligned} \tag{5.1}$$

and $u_{jk}^{\nu+1} = w_J$ is the subsequent iterate. *Hierarchical domain decomposition methods* are obtained from decompositions as presented in the preceding sections.

Theorem 5.1. *Let*

$$\mathcal{W}_0 = \mathcal{S}_j^M, \qquad \mathcal{W}_1 = \mathcal{S}_j^\Gamma, \qquad \mathcal{W}_2 = \mathcal{S}_{jk}^F$$

with $\mathcal{S}_j^M$, $\mathcal{S}_j^\Gamma$ and $\mathcal{S}_{jk}^F$ defined in (4.1), (4.8) and (3.2), respectively, and select $b_l(\cdot,\cdot) = a(\cdot,\cdot)$ for $l = 0, 1, 2$.

Then the iterates u_{jk}^ν of the resulting hierarchical domain decomposition method converge to the exact solution u_{jk} of (2.7) and satisfy the error estimate

$$\|u_{jk} - u_{jk}^{\nu+1}\|^2 \le \rho \|u_{jk} - u_{jk}^\nu\|^2, \qquad \nu = 0, \dots,$$

with $\rho < 1$ depending only on the shape regularity (2.4) of the initial partition $\mathcal{P}_0$ and on the constant C_0 in (2.6).

Proof. The proof follows from general convergence results for subspace correction methods (cf. Xu [23] or Yserentant [25]). More precisely, we can apply Theorem 5.1 in [25] with $\omega = 1$, $K_1 \preceq 1$ (cf. Propositions 3.1, 4.1) and $K_2 = 3$. □

We emphasize that Theorem 5.1 implies robust convergence with respect to arbitrary large permeability k_F and arbitrary small width ε_F of the fractures. These results are not restricted to two space dimensions. Utilizing the techniques presented in the preceding sections robust HDD algorithms with mesh-independent convergence rates can be derived for three space dimensions and for more complicated geometries like multiple crossings or fractures ending in the computational domain.

We now consider an inexact version of the above HDD algorithm replacing the exact solution of the local subproblem (5.1) for $\mathcal{W}_0 = \mathcal{S}_j^M$ by one step of a multigrid method. From now on, our considerations are limited to two space dimensions.

Successive refinement in the rock matrix gives rise to the following sequence of nested subspaces

$$\mathcal{S}_0^M \subset \cdots \subset \mathcal{S}_{j-1}^M \subset \mathcal{S}_j^M, \tag{5.2}$$

where $\mathcal{S}_l^M$ consists of all functions $v \in H_0^1(\Omega)$ such that $v|_T$ is linear for all $T \in \mathcal{T}_l$, $v|_Q$ is isoparametric bilinear for all $Q \in \mathcal{Q}_l$, and $v|_E$ is constant for all edges $E \in \mathcal{E}_l$. Recall that $\mathcal{E}_l$ and $\mathcal{N}_l$ are denoting the sets of interior edges and vertices of $P \in \mathcal{P}_l$, respectively. The underlying refinement rules imply

$$c2^{-l} \leq \operatorname{diam}(T) \leq C2^{-l} \qquad \forall T \in \mathcal{T}_l$$

with constants c, C depending only on the initial triangulation $\mathcal{T}_0$. From now on constants hidden by the shortcuts $'\asymp'$ and $'\preceq'$ may additionally depend on c, C.

The nodal interpolation $I_l : \mathcal{S}_j \to \mathcal{S}_l$ defined by

$$I_l v(p) = v(p), \qquad p \in \mathcal{N}_l, \tag{5.3}$$

gives rise to the direct splitting

$$\mathcal{S}_j^M = \mathcal{V}_0 \oplus \mathcal{V}_1 \oplus \cdots \oplus \mathcal{V}_j \tag{5.4}$$

into the subspaces

$$\mathcal{V}_0 = \mathcal{S}_0^M, \qquad \mathcal{V}_l = (I_l - I_{l-1})\mathcal{S}_j^M, \quad l = 1, \ldots, j. \tag{5.5}$$

We state a variant of the well-known stability result of Yserentant [24]. Recall the weighted L^2-norm $\|\cdot\|_{0,K}$ defined in (2.8).

Proposition 5.1. *For each $v \in \mathcal{S}_j^M$ the decomposition $v = \sum_{l=0}^j v_l$ into $v_l \in V_l$ satisfies*

$$\|v_0\|^2 + \sum_{l=1}^{j} 4^l \|v_l\|_{0,K}^2 \preceq (1+j)^2 \|v\|^2. \tag{5.6}$$

Proof. Let $v \in \mathcal{S}_j^M$. First note that

$$4^l \|(I_l - I_{l-1})v\|_{0,K}^2 \preceq \|I_l v\|^2, \quad l = 1, \ldots, j,$$

can be shown by transformation to reference elements $\hat{T}$ or $\hat{Q}$ and exploiting the equivalence of norms on 2- or 5-dimensional quotient spaces. Hence, it is sufficient to prove

$$\|I_l v\|^2 \preceq (1+j-l)\|v\|^2, \qquad l = 0, \ldots, j. \tag{5.7}$$

Obviously, (5.7) is a consequence of the local estimate

$$\|\nabla I_l v\|_{L^2(P)}^2 \preceq (1+j-l)\|\nabla v\|_{L^2(P)}^2 \qquad \forall P \in \mathcal{P}_l = \mathcal{T}_l \cup \mathcal{Q}_l$$

which is well-known for $T \in \mathcal{T}_l$ (cf. Yserentant [24]). In order to show

$$\|\nabla I_l v\|_{L^2(Q)}^2 \preceq \|\nabla v\|_{L^2(Q)}^2, \quad \forall Q \in \mathcal{Q}_l \tag{5.8}$$

we set $w = I_l v$ and let $\hat{w}$, $\hat{v}$ denote the transformed functions on the reference element $\hat{Q} = [0,1] \times [0,\varepsilon]$. As $\frac{\partial}{\partial \eta}\hat{w} = \frac{\partial}{\partial \eta}\hat{v} = 0$ the orthogonality

$$\int_0^1 \tfrac{\partial}{\partial \xi}\hat{w} \tfrac{\partial}{\partial \xi}(\hat{v} - \hat{w})\, d\xi$$

implies

$$\|\nabla\hat{v}\|^2_{L^2(\hat{Q})} = \|\nabla\hat{w}\|^2_{L^2(\hat{Q})} + \|\nabla(\hat{v}-\hat{w})\|^2_{L^2(\hat{Q})} \geq \|\nabla\hat{w}\|^2_{L^2(\hat{Q})}$$

so that (5.8) follows from Lemma 3.1. This proves the assertion. □

Next we extend a sharpened Cauchy-Schwarz inequality from classical finite element spaces to $\mathcal{S}_l^M$. Recall the weighted L^2-norm $\|\cdot\|_{0,K}$ as introduced in (2.8).

Proposition 5.2. *Let $0 \leq k < l \leq j$. Then the estimate*

$$a(v,w) \preceq \left(\tfrac{1}{\sqrt{2}}\right)^{l-k} \|v\| 2^l \|w\|_{0,K} \tag{5.9}$$

holds for all $v \in \mathcal{S}_k^M$ and $w \in \mathcal{S}_l^M$.

Proof. It is sufficient to show the local estimate

$$(\nabla v, \nabla w)_{L^2(P)} \preceq (\frac{1}{\sqrt{2}})^{l-k} \|\nabla v\|_{L^2(P)} 2^l \|w\|_{L^2(P)} \qquad \forall P \in \mathcal{T}_k \cup \mathcal{Q}_k. \tag{5.10}$$

As (5.10) is well-known for $T \in \mathcal{T}_l$ (cf., e.g., [25, p. 313]), we only have to consider the case $P = Q \in \mathcal{Q}_k$. Then the transformation to the associated reference element $\hat{Q}$ leads to

$$(\nabla v, \nabla w)_{L^2(Q)} = \varepsilon \int_0^1 \alpha(\xi)\hat{v}_\xi \hat{w}_\xi \, d\xi$$

because v and w are constant along edges $E \in \mathcal{E}_k$. Here $\alpha(\xi)$ is a quadratic polynomial with the properties

$$\alpha(\xi) \asymp 1, \quad |\alpha'(\xi)| \preceq 1 \qquad \forall \xi \in [0,1].$$

Now (5.10) follows from a one-dimensional version of the arguments used in the proof of Lemma 6.1 in [25]. □

We now present an inexact variant of HDD which requires a multigrid V-cycle and the exact solution of local problems on $\mathcal{S}_j^\Gamma$ and $\mathcal{S}_{jk}^F$ in each iteration step.

Theorem 5.2. *Let*

$$\mathcal{W}_l = \mathcal{S}_l^M, \quad l = 0,\dots,j, \qquad \mathcal{W}_{j+1} = \mathcal{S}_j^\Gamma, \qquad \mathcal{W}_{j+2} = \mathcal{S}_{jk}^F$$

with $\mathcal{S}_l^M$, $\mathcal{S}_j^\Gamma$ and $\mathcal{S}_j^F$ defined in (5.2), (4.8) and (3.2), respectively. Let $b_l(\cdot,\cdot)$, be generated by symmetric Gauß-Seidel smoothers for $l = 1,\dots,j$ and $b_l(\cdot,\cdot) = a(\cdot,\cdot)$ for $l = 0, j+1, j+2$.

Then the iterates u_{jk}^ν of the resulting hierarchical domain decomposition method converge to the exact solution u_{jk} of (2.7) and satisfy the error estimate

$$\|u_{jk} - u_{jk}^{\nu+1}\|^2 \leq \left(1 - C(1+j)^{-3}\right) \|u_{jk} - u_{jk}^\nu\|^2, \qquad \nu = 0,\dots,$$

with C depending only on the initial partition $\mathcal{P}_0$ and the constant C_0 from (2.6).

Proof. We select the direct splitting

$$\mathcal{S}_{jk} = \mathcal{V}_0 \oplus \mathcal{V}_1 \oplus \cdots \oplus \mathcal{V}_J, \quad J = j+2, \tag{5.11}$$

into the subspaces $\mathcal{V}_l \subset \mathcal{W}_l$, $l = 0,\dots,j$, defined in (5.5), $\mathcal{V}_{j+1} = \mathcal{S}_j^{\Gamma_0} \subset \mathcal{W}_{j+1}$, and $\mathcal{V}_{j+2} = \mathcal{S}_{jk}^F = \mathcal{W}_{j+2}$, defined in (4.2) and (3.2), respectively. The smoothing property (2.33) in [25] holds with $\omega = 1$ for the Gauß-Seidel smoothers $b_l(\cdot,\cdot)$, $l = 1,\dots,j$, and is trivial otherwise. The stability (5.2) in [25] of the decomposition

(5.11) with $K_1 \preceq (1+j)^2$ follows from Propositions 3.1, 4.1, and 5.1 utilizing the norm equivalence

$$b_l(v,v) \asymp 4^l \|v\|_{0,K}^2 \quad \forall v \in \mathcal{S}_l^M, \qquad l = 1,\dots,j. \tag{5.12}$$

Note that the refinement condition (2.6) is used in the proof of (5.12). Inserting (5.12) into (5.9) and using the smoothing property, we obtain the Cauchy-Schwarz-type inequality

$$a(w_k,w_l) \preceq \left(\frac{1}{\sqrt{2}}\right)^{|l-k|} b_k(w_k,w_k)^{\frac{1}{2}} b_l(w_l,w_l)^{\frac{1}{2}} \quad \forall w_k \in \mathcal{W}_k, w_l \in \mathcal{W}_l.$$

This inequality directly leads to

$$\|\sum_{l=0}^{j} w_l\|^2 \preceq \sum_{l=0}^{j} b_l(w_l,w_l) \qquad \forall w_l \in \mathcal{W}_l, \quad l = 0,\dots,j,$$

and the Cauchy-Schwarz inequality provides condition (5.7) in [25] with $K_2 \preceq 1$. Now the assertion follows from Theorem 5.4 in [25]. □

The proof of Theorem 5.2 directly applies to other smoothers satisfying (5.12). Even suitable non-symmetric smoothers like, e.g., the Gauß-Seidel method are allowed (cf. Neuss [17]).

According to Theorem 5.2 the inexact version of HDD preserves robustness with respect to ε_F and, as a consequence of the local properties of the interpolation operators I_l, with respect to k_F. On the other hand, stability of I_l is restricted to two space dimensions. Replacing interpolation by L^2-type projections, similar results can be obtained for three dimensions at the expense of robustness with respect to k_F.

The exact solution of the local subproblems associated with the fracture space $\mathcal{S}_{jk}^F$ can also be replaced by one step a multigrid method. Indeed, Lemma 3.2 provides the stability of an hierarchical splitting of the fracture space $\mathcal{S}_{jk}^F$ into subspaces of $\mathcal{S}_{jl}^F$, $l = 0,\dots,k$, of functions $v \in \mathcal{S}_{jk}^F$ which are linear on all edges $E \in \mathcal{E}_{jl} \cap \overline{\Omega}^F$. Here, $\mathcal{E}_{jl}$ denotes the edges of $P \in \mathcal{P}_{jl}$. The convergence rate of the resulting hierarchical multigrid method with line Gauß-Seidel smoother is robust for $\varepsilon \to 0$ and independent of the mesh size. For similar results we refer to Bramble and Zhang [5] and the references cited therein.

Finally note that the "robust" smoothers proposed by Gebauer et al. [10] for the multigrid solution of problems on $\mathcal{S}_j^{\overline{M}} = \mathcal{S}_j^\Gamma + \mathcal{S}_j^M$ can be interpreted in terms of a suitable multilevel splitting of $\mathcal{S}^\Gamma$.

6. Numerical results

We consider the model problem

$$\nabla \cdot (K \nabla u) = 0 \qquad \text{on} \qquad \Omega = (0,6) \times (0,6) \tag{6.1}$$

with K defined in (2.3), $u(0,y) = 2$, $u(6,y) = 1$ for $y \in [0,6]$ and homogeneous Neumann data elsewhere. Obviously, (6.1) can be written in weak form (2.2) with suitable H and ℓ. The fracture network Ω_F together with the initial partition $\mathcal{P}_0$ is shown in the left picture of Figure 2.1 for a comparatively large width $\varepsilon_F = 0.2$. Corresponding partitions for smaller ε_F are obtained by shifting the nodes lying on the interface Γ towards the centerlines of the fractures. In the limit case $\varepsilon_F = 0$ the fractures disappear and the problem reduces to the Laplace equation.



FIGURE 6.1. Robustness with respect to increasing k_F and vanishing ε_F



FIGURE 6.2. Influence of large ε_F/h_Q and small interior angles γ

In order to illustrate the robustness of the hierarchical domain decomposition method presented in Theorem 5.2, we consider the corresponding discretized problem (2.7) for $j = 6$ and $k = 2$. In the left picture of Figure 6.1 we depict the convergence rates for fixed $\varepsilon_F = 10^{-5}$ and increasing permeability k_F. More precisely, the convergence rates are approximated by

$$\rho = \frac{\|u_{jk}^{\nu_0+1} - u_{jk}^{\nu_0}\|}{\|u_{jk}^{\nu_0} - u_{jk}^{\nu_0-1}\|}$$

where ν_0 is chosen such that $\|u_{jk}^{\nu_0+1} - u_{jk}^{\nu_0}\| \leq 10^{-12}$. As expected from the theoretical findings, the convergence speed is hardly influenced by the size of the jump. The right picture shows (approximate) convergence rates for fixed $k_F = 1$ and decreasing ε_F. The convergence rates are almost the same for $10^{-9} \leq \varepsilon_F \leq 10^{-2}$. They scarcely differ from the convergence rates of classical multigrid for the reduced Laplace problem which are indicated by the horizontal line. Note that we have $C_0 \approx 40$ in condition (2.6) for $\varepsilon_F = 10^{-1}$ which explains the unsatisfying convergence speed for this value.

We now compare the convergence rates for fixed $k_F = 10^6$ and increasing number of refinement steps j. The left picture in Figure 6.2 shows that the convergence speed rapidly deteriorates for "large" $\varepsilon_F = 10^{-1}$ (upper curve) and is hardly affected for "small" $\varepsilon_F = 10^{-5}$ (lower curve). Note that $C_0 \approx 2 \cdot 10^{-3}$ in the latter case. The right picture illustrates the influence of decreasing interior angles. The length of an edge $E \in \Gamma$ of an element $Q \in \mathcal{Q}_0$ is shifted by a fixed factor s (independent of ε_F). This leads to an interior angle $\gamma \approx \arctan(\varepsilon_F/s)$ which obviously tends to zero for increasing s and small ε_F. It is interesting to see how convergence rates branch off for increasing s or, equivalently, for γ becoming too small. These two experiments complement our analysis in the sense that moderate constants in the conditions (2.4) and (2.6) also seem to be necessary for fast convergence.

Let us remark in closing that comparisons with the algebraic multigrid method by Ruge and Stüben [19] in Gebauer [9] confirm the superiority of hierarchical domain decomposition not only from a theoretical but also from a numerical point of view.

Acknowledgment . *Special thanks to R. Helmig (Uni Stuttgart), R. Hinkelmann (TU Berlin), L. Neunhäuserer (Uni Stuttgart), and S. Ochs (Uni Stuttgart) for stimulating discussions and fruitful cooperation on flow and transport in fractured porous media.*

References

[1] *T. Apel and J. Schöberl. Multigrid methods for anisotropic edge refinement.* SIAM J. Numer. Anal.*, 40:1993–2006, 2002.*

[2] *P. Bastian, Z. Chen, R.E. Ewing, R. Helmig, H. Jakobs, and V. Reichenberger. Numerical solution of multiphase flow in fractured porous media. In Z. Chen, R.E. Ewing, and Z.C. Shi, editors,* Numerical treatment of multiphase flows in porous media*, pages 50–68. Springer, 2000.*

[3] *J. Bergh and J. Löfström.* Interpolation Spaces*. Springer, Berlin, 1976.*

[4] *D. Braess. Towards algebraic multigrid for elliptic problems of second order.* Computing*, 55:379–393, 1995.*

[5] *J.H. Bramble and X. Zhang. Uniform convergence of the multigrid V-cycle for an anisotropic problem.* Math. Comp.*, 70:453–470, 2001.*

[6] *S. Brenner and L.R. Scott.* The Mathematical Theory of Finite Element Methods*. Springer, New York, 1994.*

[7] *P. Deuflhard and R. Hochmuth. Multiscale analysis of thermoregulation in the human microvascular system.* Math. Mod. Meth. in the Appl. Sci.*, 27:971–989, 2004.*

[8] *P. Deuflhard and R. Hochmuth. Multiscale analysis for the bio-heat-transfer equation.* Math. Mod. Meth. in the Appl. Sci.*, to appear.*

[9] *S. Gebauer.* Hierarchische Gebietszerlegung für die gesättigte Grundwasserströmung in Kluftaquifersystemen*. PhD thesis, Institute for Mathematics II, FU Berlin, 2004.*

[10] *S. Gebauer, L. Neunhäuserer, R. Kornhuber, S. Ochs, R. Hinkelmann, and R. Helmig. Equidimensional modelling of flow and transport processes in fractured porous systems I. In Hassanizadeh et al., editor,* Computational Methods in Water Resources'*, pages 335–342. Elsevier, 2002.*

[11] *M. Heisig, R. Lieckfeldt, G. Wittum, G. Mazurkevich, and G. Lee. Non steady-state descriptions of drug permeation through stratum corneum I: The biphasic brick-and-mortar-model.* Pharmaceutical research*, 13:421–426, 1996.*

[12] *R. Helmig.* Theorie und Numerik der Mehrphasenströmungen in geklüftet porösen Medien*. PhD thesis, Universität Hannover, 1993.*

[13] *O. Kolditz, J. de Jonge, M. Beinhorn, M. Xie, T. Kalbacher, W. Wang, S. Bauer, C.I. McDermott, R. Kaiser, and M. Kohlmeier.* ROCKFLOW

- Theory and users manual. Release 3.9. Preprint 2003-37, Center of Applied Geosciences, Geohydrology/HydroInformatics, University of Tübingen, 2003.

[14] *L. Neunhäuserer.* Diskretisierungsansätze zur Modellierung von Strömungs- und Transportprozessen in geklüftet-porösen Medien. *PhD thesis, Institut für Wasserbau, Universität Stuttgart, 2003.*

[15] *L. Neunhäuserer, S. Gebauer, S. Ochs, R. Hinkelmann, R. Kornhuber, and R. Helmig. Equidimensional modelling of flow and transport processes in fractured porous systems II. In Hassanizadeh et al., editor,* Computational Methods in Water Resources'*, pages 343–350. Elsevier, 2002.*

[16] *N. Neuss.* Homogenisierung und Mehrgitter. *PhD thesis, ICA, Universität Stuttgart, 1996.*

[17] *N. Neuss. V-cycle convergence with unsymmetric smoothers and application to an anisotropic model problem.* SIAM J.Numer.Anal.*, 35:1201–1212, 1998.*

[18] *A. Quarteroni, M. Tuveri, and A. Veneziani. Computational vascular fluid dynamics: problems, models and methods.* Comp. Vis. Sci.*, 2:163–197, 2000.*

[19] *J. Ruge and K. Stüben. Algebraic multigrid. In S.F. McCormick, editor,* Multigrid methods*, volume Vol. 3 of* Frontiers in Applied Mathematics*, pages 73–130, Philadelphia, 1987. SIAM, AMS.*

[20] *A. Silberhorn-Hemminger.* Modellierung von Kluftaquifersystemen: Geostatistische Analyse und deterministisch - stochastische Kluftgenerierung. *PhD thesis, Institut für Wasserbau, Universität Stuttgart, 2001.*

[21] *A. Ženíšek and M. Vanmaele. The interpolation theorem for narrow quadrilateral isoparametric finite elements.* Numer. Math.*, 72:123–141, 1995.*

[22] *C. Wagner. Introduction to algebraic multigrid. Course notes, Universität Heidelberg, 1998.*

[23] *J. Xu. Iterative methods by space decomposition and subspace correction.* SIAM Review*, 34:581–613, 1992.*

[24] *H. Yserentant. On the multi-level splitting of finite element spaces.* Numer. Math.*, 49:379–412, 1986.*

[25] *H. Yserentant. Old and new convergence proofs for multigrid methods.* Acta Numerica*, pages 285–326, 1993.*

Dr. Susanna Gebauer, Freie Universität Berlin, Institut für Mathematik II, Arnimallee 2-6, D - 14195 Berlin, Germany

E-mail address: `susanna.gebauer@math.fu-berlin.de`

Prof. Dr. Ralf Kornhuber, Freie Universität Berlin, Institut für Mathematik II, Arnimallee 2-6, D - 14195 Berlin, Germany

E-mail address: `kornhuber@math.fu-berlin.de`

Prof. Dr. Harry Yserentant, Technische Universität Berlin, Institut für Mathematik, Strasse des 17. Juni 136, D-10632 Berlin Germany

E-mail address: `yserentant@math.tu-berlin.de`